

Enhanced effect of temporal variation of the fine structure constant in diatomic molecules

V.V. Flambaum

School of Physics, The University of New South Wales, Sydney NSW 2052, Australia

(Dated: February 2, 2008)

We show that the relative effect of variation of the fine structure constant in microwave transitions between very close and narrow rotational-hyperfine levels may be enhanced 2-3 orders of magnitude in diatomic molecules with unpaired electrons like LaS, LaO, LuS, LuO, YbF and similar molecular ions. The enhancement is result of cancellation between the hyperfine and rotational intervals.

PACS numbers: 06.20.Jr , 33.20.Bx , 31.30.Jv

The possible variation of the fundamental constants of nature is currently a very popular research topic. There are hints for the variation of the fine structure constant, $\alpha = e^2/\hbar c$, strength constant of the strong interaction and masses in Big Bang nucleosynthesis, quasar absorption spectra and Oklo natural nuclear reactor data (see e.g.[1]) . However, a majority of publications report only limits on possible variations (see e.g. reviews [2, 3]). A very sensitive method to study the variation in a laboratory consists of the comparison of different optical and microwave atomic clocks (see recent measurements in [4, 5, 6, 7, 8, 9, 10]). An enhancement of the relative effect of α variation can be obtained in transition between the almost degenerate levels in Dy atom [11]. These levels move in opposite directions if α varies. An experiment is currently underway to place limits on α variation using this transition [12]. Unfortunately, one of the levels has quite a large linewidth and this limits the accuracy.

In this paper I would like to note that very close narrow levels exist in diatomic molecules. In the work [13] we suggested the use of such levels to enhance the effects of parity violation. Below I will show that there is also an enhancement of α variation in transitions between such levels.

Consider, for example, the molecule $^{139}\text{La}^{32}\text{S}$ (or LaO, LuO, LuS, YbF, etc, or similar molecular ions with unpaired valence electrons). The electron ground state of this molecule is $^2\Sigma$. The rotational-hyperfine levels are characterized by the total angular momentum $\mathbf{J} = \mathbf{L} + \mathbf{F}$, rotational angular momentum L , $\mathbf{F} = \mathbf{I} + \mathbf{S}$, nuclear spin I and electron spin S . The interval between the rotational-hyperfine levels $|L = 1, F = 4, J >$ and $|L = 2, F = 3, J' >$ is close to zero within the experimental accuracy (about 0.01 cm^{-1} , see all data in [14]). This is a result of the cancellation between the rotational interval $B[L_1(L_1 + 1) - L_2(L_2 + 1)]$ ($B=0.11693 \text{ cm}^{-1}$) and the hyperfine interval $4b$ ($4b=0.47 \text{ cm}^{-1}$); the transition frequency is $\omega = 4b - 4B = 0.002 \pm 0.01 \text{ cm}^{-1}$.

The hyperfine structure constant can be presented in the following form

$$b = \text{const} \times \left[\frac{m_e e^4}{\hbar^2} \right] [\alpha^2 F_{\text{rel}}(Z\alpha)] \left[\mu \frac{m_e}{m_p} \right] \quad (1)$$

The factor in the first bracket is the atomic unit of en-

ergy. The second “electromagnetic” bracket determines the dependence on α . An approximate expression for the relativistic correction factor (Casimir factor) for s-wave electron is the following

$$F_{\text{rel}} = \frac{3}{\gamma(4\gamma^2 - 1)} \quad (2)$$

where $\gamma = \sqrt{1 - (Z\alpha)^2}$, Z is the nuclear charge. Variation of α leads to the following variation of F_{rel} [4]:

$$\frac{\delta F_{\text{rel}}}{F_{\text{rel}}} = K \frac{\delta \alpha}{\alpha} \quad (3)$$

$$K = \frac{(Z\alpha)^2(12\gamma^2 - 1)}{\gamma^2(4\gamma^2 - 1)} \quad (4)$$

More accurate numerical many-body calculations [11] of the dependence of the hyperfine structure on α have shown that the coefficient K is slightly larger than that given by this formula. For Cs ($Z=55$) $K=0.83$ (instead of 0.74), for Hg^+ $K=2.28$ (instead of 2.18). The last bracket in eq. (1) contains the dimensionless nuclear magnetic moment μ in nuclear magnetons (the nuclear magnetic moment $M = \mu \frac{e\hbar}{2m_p c}$), electron mass m_e and proton mass m_p . The dependence of μ on the dimensionless strong interaction parameter $X = m_q/\Lambda_{QCD}$ has been calculated in Ref. [15, 16]. Here m_q is the quark mass and Λ_{QCD} is the strong interaction scale.

The rotational constant can be presented in the following form

$$B = \text{const} \times \left[\frac{m_e e^4}{\hbar^2} \right] \left[\frac{m_e}{m_p} \right] \quad (5)$$

Using eqs. (1,5) we obtain the frequency

$$\omega = \text{const} \times \left[\frac{m_e e^4}{\hbar^2} \right] \left[\frac{m_e}{m_p} \right] [\alpha^2 F_{\text{rel}}(Z\alpha)\mu - \text{const}] \quad (6)$$

We are interested in the case when two terms in the last bracket nearly cancel each other ($\omega = 4b - 4B \ll 4b$). In this case ω is very sensitive to the variation of α and μ . The relative variation of the transition frequency is then approximately equal to

$$\frac{\delta \omega}{\omega} \approx \frac{4b}{\omega} \left[(2 + K) \frac{\delta \alpha}{\alpha} + \beta \frac{\delta X}{X} \right] \quad (7)$$

For La ($Z=57$) $K = 0.9$ and $\beta \sim 0.01$ [16]. For a numerical estimate we can assume the central value for the frequency, $\omega = 4b - 4B = 0.002 \text{ cm}^{-1}$. Then we have

$$\frac{\delta\omega}{\omega} \sim 600 \frac{\delta\alpha}{\alpha} \quad (8)$$

The accurate value of the coefficient before $\delta\alpha/\alpha$ can be obtained after the frequency ω is measured. Thus in molecular experiments one can have a two to three orders of magnitude enhancement of the relative effect of α variation. Present laboratory sensitivity to the variation of α has reached $2 \cdot 10^{-15}$ per year (see review [3] and measurements [5, 6, 7, 8, 9, 10]). Thus, the required frequency stability should be about

$$\frac{\delta\omega}{\omega} \sim 10^{-12} \quad (9)$$

For comparison, the microwave clocks have stability

about 10^{-16} (see e.g. [6]). It is also instructive to present an absolute shift of the frequency:

$$\delta\omega \approx (2 + K)4b \frac{\delta\alpha}{\alpha} \approx 4 \cdot 10^{10} \frac{\delta\alpha}{\alpha} \text{ Hz} \quad (10)$$

This corresponds to $\delta\omega \sim 10^{-4}$ Hz per year. The natural width of rotational levels in the ground state is small. Therefore, the actual width and sensitivity to the variation of α in molecules will depend on the particular experimental arrangement which is beyond the scope of the present paper. Here we only would like to note that it was suggested in Ref. [17] to trap molecular ions for the measurements of the electron electric dipole moment. A similar trapping scheme may possibly be used to study the variation of α in molecular ions.

This work is supported by the Australian Research Council.

-
- [1] M.T.Murphy, J.K. Webb, V.V. Flambaum. Mon. Not. R. Astron. Soc. **345**, 609-638 (2003). V.F. Dmitriev, V.V. Flambaum, J.K. Webb, Phys. Rev. D **69**, 063506 (2004). S K Lamoreaux and J R Torgerson *Phys. Rev. D* **69** 121701 (2004). A. Ivanchik, P. Petitjean, D. Aracil, R. Strianand, H. Chand, C. Ledoux, P. Boisse *Astron. Astrophys.* **440**, 45 (2005).
 - [2] J-P. Uzan, Rev. Mod. Phys. **75**, 403 (2003);
 - [3] S.G. Karshenboim, V.V. Flambaum, E. Peik. Atomic clocks and constraints on variations of fundamental constants. In "Handbook of atomic, molecular and optical physics" (Ed. G.W.F. Drake, Springer 2005), Chapter 30, pp.455-463; arXiv:physics/0410074.
 - [4] J.D. Prestage, R.L. Tjoelker, and L. Maleki. Phys. Rev. Lett **74**, 3511 (1995).
 - [5] H. Marion *et al.*, Phys. Rev. Lett. **90**, 150801 (2003).
 - [6] S. Bize *et al.*, arXiv:physics/0502117.
 - [7] E. Peik, B. Lipphardt, H. Schnatz, T. Schneider, Chr. Tamm, S.G. Karshenboim. Phys. Rev. Lett. **93** 170801 (2004); arXiv:physics/0402132 .
 - [8] S. Bize *et al.*, Phys. Rev. Lett. **90**, 150802 (2003).
 - [9] M. Fischer *et al.* arXiv:physics/0311128.
 - [10] E. Peik, B. Lipphardt, H. Schnatz, T. Schneider, Chr. Tamm, S.G. Karshenboim, arXiv:physics/0504101.
 - [11] V.A. Dzuba, V.V. Flambaum, J.K. Webb. Phys. Rev. A **59**, 230 (1999).
 - [12] A T Nguyen, D Budker, S K Lamoreaux and J R Torgerson *Phys. Rev. A*. **69**, 022105 (2004).
 - [13] V.V. Flambaum, I.B. Khriplovich. Phys. Lett. A **110**, 121 (1985).
 - [14] K.P. Huber and G. Herzberg. Constants of diatomic molecules (Van Nostrand Reinhold Company, New York, 1979; Mir, Moscow, 1984).
 - [15] V.V. Flambaum, physics/0309107. V.V. Flambaum, D.B. Leinweber, A.W. Thomas, R.D. Young, Phys. Rev. D **69**, 115006 (2004).
 - [16] V.V. Flambaum, A.F. Tedesco, arXiv:nucl-th/0601050 .
 - [17] E. Stutz and E. Cornell., Bull. Amer. Phys. Soc. **49**, 76 (2004).